09/856886 .072601

JC10 Rec'd PCT/PTO 25 MAY 2001

FORM PTO-1390 (REV. 10-2000)	DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY DOCKET NUMBER IT0013P
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 09/856886
INTERNATIONAL APPLICATION NO. PCT/EP99/09207	INTERNATIONAL FILING DATE 24/11/1999	PRIORITY DATE CLAIMED 26/11/1998

TITLE OF INVENTION

PHARMACEUTICAL COMPOUNDS FOR THE INHIBITION OF HEPATITIS C VIRUS NS3 PROTEASE

APPLICANT(S) FOR DO/EO/US

ANTONELLO PESSI, PAOLA INGALLINELLA, ELISABETTA BIANCHI

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. [X] This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. [] This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. [] This is an express request to begin national examination procedures [35 U.S.C. 371(f)] at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. [X] A proper Demand for International Preliminary Examination was made and the US was elected by the expiration of the 19th month from the earliest claimed priority date (PCT Article 31).
5. [X] A copy of the International Application as filed [35 U.S.C. 371(c)(2)].
 a. [X] is attached hereto (required only if not communicated by the International Bureau).
 b. [] has been communicated by the International Bureau.
 c. [] is not required, as the application was filed in the United States Receiving Office (RO/US).
6. [] An English language translation of the International Application as filed [35 U.S.C. 371(c)(2)].
7. [] Amendments to the claims of the International Application under PCT Article 19 [35 U.S.C. 371(c)(3)].
 a. [] are attached hereto (required only if not communicated by the International Bureau).
 b. [] have been communicated by the International Bureau.
 c. [] have not been made; however, the time limit for making such amendments has NOT expired.
 d. [] have not been made and will not be made.
8. [] An English language translation of the amendments to the claims under PCT Article 19 [35 U.S.C. 371(c)(3)].
9. [] An oath or declaration of the inventor(s) [35 U.S.C. 371(c)(4)].
10. [] An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 [35 U.S.C. 371(c)(5)].

Items 11 to 16 below concern other document(s) or information included:

11. [] An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. [] An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. [X] A **FIRST** preliminary amendment.
 [] A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. [] A substitute specification.
15. [] A change of power of attorney and/or address letter.
16. [] Other items or information:

EXPRESS MAIL CERTIFICATE
DATE OF DEPOSIT MAY 25 2001
EXPRESS MAIL NO. EM 230302696 US
I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL "POST OFFICE TO ADRESSEE" BEFORE 5 P.M. ON THE ABOVE DATE IN AN ENVELOPE ADDRESSED TO ASSISTANT COMMISSIONER FOR PATENTS, WASHINGTON, D.C., 20231
MAILED BY Kathy Kleinhauser
DATE 5-25-01

Computer generated form "TransLetter (DO-EO-US)" (PCT Folder), Merck & Co., Inc., 10/3/00

09856886 09/856886

PATENT

JC18 Rec'd PCT/PTO 25 MAY 2001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants:	Pessi, et al.
Serial No.:	Case No.: IT0013P
Filed:	
For:	PHARMACEUTICAL COMPOUNDS FOR THE INHIBITION OF HEPATITIS C VIRUS NS3 PROTEASE

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Please amend the application filed herewith as provided below.

In the Claims:

Please cancel claim 13 without prejudice to future prosecution.

Please amend the claims by replacing and adding claims as follows:

3. (Amended Once) The compound of claim 1, a pharmaceutically acceptable salt or derivative thereof which is N-terminally acylated.

4. (Amended Once) A compound, salt or derivative according to claim 1 wherein A'-B'-C'-D' is a tetrapeptide of formula:

Pro-B'-C'-Leu

wherein B' and C' are as defined in claim 1.

EXPRESS MAIL CERTIFICATE
DATE OF DEPOSIT MAY 25 2001
EXPRESS MAIL NO. EM 230307696 US
I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL "POST OFFICE TO ADDRESSEE" BEFORE 5 P.M. ON THE ABOVE DATE IN AN ENVELOPE ADDRESSED TO ASSISTANT COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231.
MAILED BY Kathy L. Lufkowski
DATE 5-25-01

5. (Amended Once) A compound, salt or derivative according to claim 4 wherein B' is selected from: β-cyclohexylalanine, phenylglycine, homophenylalanine, norleucine, leucine, methionine, norvaline and β-cyclopropylalanine.

7. (Amended Once) A compound, salt or derivative according to claim 5 wherein C' is selected from: aspartic acid, glutamic acid, γ-carboxyglutamic acid, glutamine, asparagine, hydroxyproline, N-β-Aloc- diaminobutyric acid, thiazolylalanine, methionine sulfoxide, pyridylalanine and serine.

9. (Amended Once) A compound, salt or derivative according to claim 4, wherein the combination of amino acids B'C' is selected from:

Cha-Ser
Cha-Asp
Nle-Asp
Hof-Asp
Phg-Asp
Cha-Gln
Nle-Gln
Hof-Gln
Cha-Hyp
Nle-Hyp
Hof-Hyp
Nle-Ser.

10. (Amended Once) A compound, salt, or derivative according to claim 9 wherein Pep-OH is capable of binding HCV NS3 protease, in the absence of the C-terminal residues A'-B'-C'-D', and has an IC_{50} below 100μM in an inhibition assay.

11. (Amended Once) A compound, salt, or derivative according to claim 9 wherein Pep is a hexa-, penta- or tetra-peptide having formula (II) below:

F-E-D-C-B-A

wherein: A is an amino acid or amino acid analogue having an aliphatic side chain of from 1 to 6 carbon atoms;

B is an amino acid or analogue having a non-polar, acidic, or polar but uncharged side group;

C is an amino acid or amino acid analogue having a non-polar or acidic side chain;

D is an amino acid or amino acid analogue having a hydrophobic side group;

E together with F may be absent, but if present is an amino acid or amino acid analogue having an acidic side chain, non-polar side chain or polar, but uncharged side chain, or is a dicarboxylic acid containing up to 6 carbon atoms and lacking the amino group of acidic amino acids;

and F may be absent (either by itself, or together with E) but when present is an amino acid or analogue having an acidic side chain or is a dicarboxylic acid containing up to 6 carbon atoms.

14. (Amended Once) A pharmaceutical composition comprising a compound, salt or derivative according to claim 1 and a pharmaceutically acceptable excipient, diluent or carrier.

15. (Amended Once) Use of a compound, salt or derivative according to claim 1 in the manufacture of a medicament for the treatment or prevention of hepatitis C or a related condition.

16. (Amended Once) A method of inhibiting HCV NS3 protease activity, and/or of treating or preventing hepatitis C or a related condition, comprising administering to a human or mammalian subject suffering from the condition a therapeutically or prophylactically effective amount of a composition according to claim 14.

17. (New) The composition of claim 14, wherein

A' is proline,

B' is selected from the group consisting of: β-cyclohexylalanine, phenylglycine, homophenylalanine, norleucine, leucine, methionine, norvaline and β-cyclopropylalanine;

C' is selected from the group consisting of aspartic acid, glutamic acid, γ-carboxyglutamic acid, glutamine, asparagine, hydroxyproline, N-β-Aloc- diaminobutyric acid, thiazolylalanine, methionine sulfoxide, pyridylalanine and serine,

D' is Leu,

or a pharmaceutically acceptable salt or derivative thereof.

18. (New) The composition of claim 17, wherein B' is either phenylglycine or β-cyclopropylalanine, or a pharmaceutically acceptable salt or derivative thereof.

19. (New) The composition of claim 17, wherein C' is aspartic acid, or a pharmaceutically acceptable salt or derivative thereof.

20. (New) The composition of claim 14, wherein

A' is proline,

D' is leucine,

B'C' is a combination selected from the group consisting of: Cha-Ser, Cha-Asp, Nle-Asp, Hof-Asp, Phg-Asp, Cha-Gln, Nle-Gln, Hof-Gln, Cha-Hyp, Nle-Hyp, Hof-Hyp, and Nle-Ser,

or a pharmaceutically acceptable salt or derivative thereof.

21. (New) The composition of claim 20, wherein Pep is a hexa-, penta- or tetra-peptide having formula (II) below:

F-E-D-C-B-A

A is selected from: cysteine, aminobutyric acid, di- and tri-fluoro aminobutyric acid, norvaline, allylglycine and alanine;

B is selected from: glutamic acid, aspartic acid, glycine, methyl glycine, 2-amino butyric acid, alanine, isoleucine, valine, leucine, cysteine, naphthylalanine and β-cyclohexylalanine;

C is selected from: glutamic acid, aspartic acid, glycine, methyl glycine, 2-amino butyric acid, alanine, isoleucine, valine, leucine, cysteine, naphthylalanine and β -cyclohexylalanine;

D is selected from: methionine, isoleucine, leucine, norleucine, valine, methylvaline, phenylglycine, diphenylalanine, tyrosine, thienylalanine, and chlorophenylalanine;

E is selected from: glutamic acid, aspartic acid, phenylalanine, diphenylalanine, isoleucine, valine, tyrosine, 4-nitrophenylalanine, glutaric acid and succinic acid;

and F is selected from: aspartic acid, glutamic acid, glutaric acid and succinic acid

or a pharmaceutically acceptable salt or derivative thereof.

22. (New) A method of inhibiting HCV NS3 protease activity, and/or of treating or preventing hepatitis C or a related condition, comprising administering to a human or mammalian subject suffering from the condition a therapeutically or prophylactically effective amount of a compound according to claim 1.

Remarks

The present amendment cancels claim 13, amends claims 3, 4, 5, 7, 9, 10, 11, 14, 15, and 16, and adds claims 17, 18, 19, 20, 21, 22. The cancellation of claim 13 and the changes to the claims are without prejudice to future prosecution.

The changes to the claims are being made to eliminate the use of a multiple dependent claim format, to insert some new claims previously described using a multiple dependent claim format, and to correct typographically errors in claims 9 and 10. The changes to the claims do not narrow the claimed invention, but rather correspond to originally filed multiple dependent claims.

To save on filing costs, the full range of claims eliminated by removing multiple dependencies were not reintroduced. Applicant reserves the right to reintroduce such claims at a later time.

Accordingly the claims are in condition for allowance. Please charge deposit account 13-2755 for fees due in connection with this amendment. If any time extensions are needed for the timely filing of the present amendment, Applicants petition for such extensions and authorize the charging of deposit account 13-2755 for the appropriate fees.

Respectfully submitted,

By _____

Sheldon O. Heber
Reg. No. 38,179
Attorney for Applicants

Merck & Co., Inc.
P.O. Box 2000
Rahway, NJ 07065-0907
(732) 594-1958

Case No.: IT0013P

Version with Markings to Show Changes Made

Changes to the Claims:

3. (Amended Once) The compound of claim 1 [or claim 2], a pharmaceutically acceptable salt or derivative thereof which is N-terminally acylated.

4. (Amended Once) A compound, salt or derivative according to [~~any one of claims~~] claim 1 [~~to 3~~] wherein A'-B'-C'-D' is a tetrapeptide of formula:

Pro-B'-C'-Leu

wherein B' and C' are as defined in claim 1.

5. (Amended Once) A compound, salt or derivative according to [~~any one of the previous claims~~] claim 4 wherein B' is selected from: β-cyclohexylalanine, phenylglycine, homophenylalanine, norleucine, leucine, methionine, norvaline and β-cyclopropylalanine.

7. (Amended Once) A compound, salt or derivative according to [~~any one of the previous claims~~] claim 5 wherein C' is selected from: aspartic acid, glutamic acid, γ-carboxyglutamic acid, glutamine, asparagine, hydroxyproline, N-β-Aloc- diaminobutyric acid, thiazolylalanine, methionine sulfoxide, pyridylalanine and serine.

9. (Amended Once) A compound, salt or derivative according to [~~any one of the preceding claims~~] claim 4, wherein the combination of amino acids B'C' is selected from:

Cha-Ser

Cha-Asp

Nle-Asp

Hof-Asp

Phg-Asp

Cha-Gln

Nle-Gln

Hof-Gln

Cha-Hyp

Nle-Hyp

Hof-Hyp

Nle-Ser.

10. (Amended Once) A compound, salt, or derivative according to [~~any one of the preceding claims~~] claim 9 wherein Pep-OH is capable of binding HCV NS3 protease, in the absence of the C-terminal residues A'-B'-C'-D', and has an IC_{50} below 100μM in an inhibition assay.

11. (Amended Once) A compound, salt, or derivative according to [~~any one of the preceding claims~~] claim 9 wherein Pep is a hexa-, penta- or tetra-peptide having formula (II) below:

F-E-D-C-B-A

wherein: A is an amino acid or amino acid analogue having an aliphatic side chain of [~~form~~] from 1 to 6 carbon atoms;

B is an amino acid or analogue having a non-polar, acidic, or polar but uncharged side group;

C is an amino acid or amino acid analogue having a non-polar or acidic side chain;

D is an amino acid or amino acid analogue having a hydrophobic side group;

E together with F may be absent, but if present is an amino acid or amino acid analogue having an acidic side chain, non-polar side chain or polar, but uncharged side chain, or is a dicarboxylic acid containing up to 6 carbon atoms and lacking the amino group of acidic amino acids;

and F may be absent (either by itself, or together with E) but when present is an amino acid or analogue having an acidic side chain or is a dicarboxylic acid containing up to 6 carbon atoms.

14. (Amended Once) A pharmaceutical composition comprising a compound, salt or derivative according to [~~any one of the preceding claims~~] claim 1 and a pharmaceutically acceptable excipient, diluent or carrier.

15. (Amended Once) Use of a compound, salt or derivative according to [~~any one of the preceding claims~~] claim 1 in the manufacture of a medicament for the treatment or prevention of hepatitis C or a related condition.

16. (Amended Once) A method of inhibiting HCV NS3 protease activity, and/or of treating or preventing hepatitis C or a related condition, comprising administering to a human or mammalian subject suffering from the condition a therapeutically or prophylactically effective amount of a composition according to claim 14[~~, or of a compound of any one of claims 1 to 12~~].

09856886 072601
09/856886
JC18 Rec'd PCT/PTO 2 5 MAY 2001

PHARMACEUTICAL COMPOUNDS FOR THE INHIBITION OF HEPATITIS C VIRUS NS3 PROTEASE

Technical Field

This invention relates to compounds which can act as inhibitors of the hepatitis C virus (HCV) NS3 protease, to uses of such compounds and to their preparation.

Background Art

The hepatitis C virus (HCV) is the major causative agent of parenterally-transmitted and sporadic non-A, non-B hepatitis (NANB-H). Some 1% of the human population of the planet is believed to be affected. Infection by the virus can result in chronic hepatitis and cirrhosis of the liver, and may lead to hepatocellular carcinoma. Currently no vaccine nor established therapy exists, although partial success has been achieved in a minority of cases by treatment with recombinant interferon-α, either alone or in combination with ribavirin. There is therefore a pressing need for new and broadly-effective therapeutics.

Several virally-encoded enzymes are putative targets for therapeutic intervention, including a metalloprotease (NS2-3), a serine protease (NS3), a helicase (NS3), and an RNA-dependent RNA polymerase (NS5B). The NS3 protease

is located in the N-terminal domain of the NS3 protein, and is considered a prime drug target since it is responsible for an intramolecular cleavage at the NS3/4A site and for downstream intermolecular processing at the NS4A/4B, NS4B/5A and NS5A/5B junctions.

Previous research has identified classes of peptides, in particular hexapeptides, showing degrees of activity in inhibiting the NS3 protease. The aim of the present invention is to provide further compounds which exhibit similar, and if possible improved, activity.

According to the nomenclature of Schechter & Berger (1967, Biochem. Biophys. Res. Commun. 27, 157-162) cleavage sites in substrates for the NS3 protease are designated P6-P5-P4-P3-P2-P1...P1'-P2'-P3'-P4'-, with each P representing an amino acid, and the scissile bond lying between P1 and P1'. Corresponding binding sites on the enzyme are indicated as S6-S5-S4-S3-S2-S1...S1'-S2'-S3'-S4'.

The present applicant has previously disclosed so called product inhibitors which are based on the P-region of the natural cleavage sites and which have been optimised to low nanomolar potency ((1998) Biochemistry 37: 8899-8905 and (1998) Biochemistry 37: 8906-8914). These inhibitors extract much of their binding energy

from the C-terminal carboxylate, the remaining interactions with NS3 being similar to the ones used by the natural substrates, including binding in the S_1 pocket and the prominent electrostatic interaction of the P6-P5 acidic couple.

At variance with the P region, the P′ region of the substrate, while being important for catalysis, does not influence significantly ground-state binding to the enzyme as expressed by the Km value. In other words, binding energy released by the substrate interaction with the enzyme to form an initial non-covalent complex is essentially due to the interaction of the residues of the P region; the P′region residues contribute to a lesser extent to the binding energy. Accordingly, peptides based on the P′region of the natural substrates (spanning residues P_1'-P_{10}') do not inhibit NS3 to any significant extent. This notwithstanding, inspection of the crystal structure of NS3 with or without 4A (and more recently of the NMR structure of NS3) shows the presence of binding pockets in the S′ region which might be exploited for the binding of active-site directed inhibitors. S′-binding ligands would therefore display a range of interactions with the enzyme different from the ones used by the substrate, and represent a novel class of NS3 inhibitors.

Landro et al in (1997) Biochemistry 36, 9340-9348 synthesized certain non-cleavable decapeptides based on the NS5A/5B cleavage site by substituting the P_1' serine by a bulky cyclic aromatic (tetrahydroisoquinoline-3-carboxylic acid) or smaller cyclic alkyl compound (proline or pipecolinic acid). They then investigated the interaction of these decapeptides with the substrate binding site of NS3 either in the presence or absence of NS4A cofactor. By looking at the effect of truncation at either the P or P' side of the molecule they concluded that most of the binding energy of the decapeptide is due to interactions with NS3-NS4A complex on the P side of the molecule. Truncation on the P' side produced a relatively large effect in the presence of NS4A cofactor, but less when NS4A was absent. They concluded that the P4' substrate Tyr residue present in their molecules was in close proximity, or in direct contact with NS4A and that this residue contributes significantly to binding in the presence of NS4A.

The present inventors have developed inhibitors which are more powerful than those described by Landro et al because they have better binding on their P' side. In other words, the inhibitors take advantage of binding to the S' region in addition to binding to the S-region of NS3. By varying the P' amino acid residues, the present

inventors have shown that the binding energy which may be extracted from S′-region binding is substantial, since inhibitors with optimised and non-optimised P′-regions differ in potency > 1000-fold. Since no activity was present in any of the peptides corresponding to the isolated P′-region, optimisation of an S′-binding fragment was pursued in the context of non-cleavable decapeptides spanning P_6-P_4'.

The inventors found that, by replacing Landro's P4′ Tyr residue by leucine the effectiveness of the decapeptides as NS3 protease inhibitors could be enhanced. Although it had been previously shown that leucine in position P4′ is better than tyrosine in a decapeptide substrate cleavable by NS3 (Urbani et al (1997) J. Biol. Chem 272, 9204-9209), this is the first showing that the same applies to decapeptide inhibitors which are not cleaved under the influence of the enzyme. By optimising the P4′ residue and then the P2′-P3′ fragment and using these together with an optimised P region the inventors have arrived at oligopeptides which show potency in the low nanomolar-subnanomolar range.

Disclosure of the Invention

According to a first aspect of the present invention there is provided a compound having the formula (I)

(written from N-terminus to C-terminus):

Pep-A′-B′-C′-D′ (I)

wherein "Pep" is a peptide or peptide analogue capable of binding to HCV NS3 protease; in particular, it is capable of binding in the S-region of the protease;

A′ is proline which is optionally substituted, for instance with one to three substituent groups;

B′ is an amino acid or amino acid analogue having a non polar side chain. Preferably, the side chain is an alkyl, aryl or aralkyl group containing 3 to 10, particularly 4 to 8 carbon atoms;

C′ is an amino acid or amino acid analogue having a polar side group. Examples of polar side group may contain between 2 and 10, preferably 2 to 6 carbon atoms;

D′ is leucine, or less preferably another amino acid with a non-polar aliphatic side chain, such as valine, isoleucine, norleucine or methionine. Alternatively, it is a short peptide or peptide analogue having one of these amino acids, especially leucine at its N terminus. The short peptide or peptide analogue may, for instance

comprise 2 to 6, preferably 2 to 4 amino acids or amino acid analogues.

As used herein, the term "amino acid analogue" includes organic compounds containing an amino and a carboxylic acid group, for instance arranged α- to each other, and which do not necessarily occur in nature.

The Pep-A bond of the compound of formula (I) is substantially uncleavable by HCV NS3 protease. For instance, it is preferable that no cleavage be detectable using the assay described below under the heading "Substrate Assay".

Pharmaceutically acceptable salts of the compound of formula (I), as well as derivatives, such as esters are within the scope of the present invention.

Preferably, the compound of formula (I) is N-terminally acylated, especially acetylated, although other derivatives of the N-terminus are also possible, for instance N-terminal sulphoxide, sulphonamide, urethane or urea derivatives.

Preferably, the compound of formula (I) is C-terminally amidated. However, the C-terminus may be an

underivatised carboxylic acid group. Alternatively, other C-terminal groups may be present.

Assuming no substitution of the proline residue at A' is present, then a preferred C-terminal portion of the compound of formula I is:

Pro-B'-C'-Leu

possibly with a short C terminal extension at Leu.

Preferred examples of the amino acid, or analogue, B' for inclusion in compounds of the first aspect of the invention, include:

β-cyclohexylalanine, phenylglycine, homophenylalanine and norleucine; other possibilities, though less preferred, are leucine, methionine, norvaline, and β-cyclopropylalanine. Of all these, cyclohexylalanine and phenyl glycine are most preferred.

Examples of the amino acid or analogue, C' include aspartic acid, glutamic acid, γ-carboxyglutamic acid, glutamine, asparagine, and hydroxyproline. Slightly less preferred are N-β-Aloc-diaminobutyric acid, thiazolylalanine, methionine sulfoxide, pyridylalanine

and serine. Of all of these aspartic acid is most preferred.

The following combinations of amino acid residues at B′ and C′ are preferred, of which the combination of cyclohexylalanine and aspartic acid is especially preferred.

TABLE 1

B′	C′
Cha	Ser
Cha	Asp
Nle	Asp
Hof	Asp
Phg	Asp
Cha	Gln
Nle	Gln
Hof	Gln
Cha	Hyp
Nle	Hyp
Hof	Hyp
Nle	Ser

Notes: Cha = β-cyclohexylalanine.
Nle = norleucine.
Phg = phenylglycine.
Hof = homophenylalanine.
Hyp = hydroxyproline.

When the residue D′ is leucine (or other amino acid)

with a small peptide as C-terminal extension the peptide may be chosen by comparison with the corresponding P′ portion of natural substrates.

The residues A′,B′,C′ and D′ may have D- or L-stereochemistry, although L-stereochemistry is, in general, preferred for each.

As regards the Pep part of the compound of formula (I) this is particularly preferably a peptide or peptide analogue capable of binding to HCV NS3 protease, even in the absence of the C-terminal residues A′-B′-C′-D′, for instance when Pep carries just a carboxylic acid group at the C terminus. For example, when tested in the inhibition assay described below the fragment Pep-OH preferably has an IC_{50} below 100μM, e.g. below 20μM, particularly below 10μM and, optimally, of less than 1μM. Preferably, Pep is a hexa-, penta- or tetra peptide having formula II below:

F-E-D-C-B-A (II)

wherein: A is an amino acid or amino acid analogue having a relatively small (C_1-C_6) aliphatic side chain. Possible choices for this group include cysteine, aminobutyric acid (Abu) (including di- and tri-fluoro

Abu), norvaline, allylglycine and alanine, any of which may be N-methylated. Of these, cysteine and the fluorinated aminobutyric acids are preferred choices for A.

B is an amino acid or analogue having a non-polar or acidic side chain. Some amino acids having polar but uncharged side groups may also be suitable. Examples of suitable amino acids include glutamic and aspartic acid, glycine and methyl glycine, 2-amino butyric acid, alanine, isoleucine, valine, leucine, cysteine, naphthylalanine and β-cyclohexylalanine. Of these, cyclohexylalanine is particularly preferred.

C is an amino acid or amino acid analogue having a non-polar or acidic side chain. For instance, the examples of such amino acids given above for B apply also to C. In this case isoleucine and glutamic acid are particularly preferred.

D is usually an amino acid or amino acid analogue having a hydrophobic side group such as methionine, isoleucine, leucine, norleucine, valine, methyl valine, phenylglycine or, diphenylalanine. Among these leucine and, particularly, diphenylalanine are preferred. Some polar amino acids which include hydrophobic portions,

such as tyrosine, thienylalanine, and chlorophenylalanine may be suitable.

E together with F may be absent, but if present is generally an amino acid or amino acid analogue having an acidic side chain. Preferred examples are glutamic and aspartic acid, with the former being preferred. E may, alternatively, be an amino acid or analogue having a non-polar, or polar but uncharged side chain. Of the non-polar amino acids, phenylalanine, diphenylalanine, isoleucine and valine are preferred, especially the D-enantiomers. Among the polar amino acids suitable examples are tyrosine and 4-nitrophenylalanine. Alternatively, where F is absent (see below), E may be a dicarboxylic acid containing up to 6 carbon atoms and lacking the amino group of acidic amino acids. Suitable examples are glutaric and succinic acid.

F may be absent (either by itself, or together with E), but when present is an amino acid or analogue having an acidic side chain. Aspartic acid is preferred, although glutamic acid is another possibility. Like E, F may also be a dicarboxylic acid containing up to 6 carbon atoms, and lacking the amino group of acidic amino acids. Examples are glutaric and succinic acid.

Of residues E and F preferably at least E is present. Particularly preferably both are present.

The amino acids and analogues A-F may be either L- or D- enantiomers though L- is generally preferred for all residues. In some cases it may be beneficial for one or other of the residues to be D- while the rest are L-. In particular it may be advantageous for E to be D-glu.

Preferred examples of the peptide "Pep" are listed below in Tables 2 and 3 together with their IC_{50} values when unextended at the C-terminus. Except for the compounds having a succinyl residue at the N-terminus, all compounds tested were N-acetylated at the N-terminus.

TABLE 2

Exp No.	Sequence	IC_{50} (μM)
1	Asp Glu Met Glu Glu Cys	1.0
2	Asp Glu Met Glu Glu D-Cys	4.0
3	Asp Glu Met Glu Glu Abu	5.8
4	Met Glu Glu Cys	150.0
5	Glu Met Glu Glu Cys	21.0
6	Glu Asp Val Val Cys Cys	5.3
7	Glu Asp Val Val Abu Cys	2.8
8	Asp Glu Val Val Cys Cys	2.1
9	Glu Asp Val Val Gly Cys	20.0
10	Asp Glu Met Glu Glu Alg	12.0
11	Glu Asp Val Val MGly Cys	21.0
12	Glu Asp MVal Val Abu Cys	1.3
13	GluS Met Glu Glu Cys	1.3
14	AsGlu Met Glu Glu Cys	0.6
15	Asp Glu Met Glu Leu Cys	1.1
16	Asp Glu Met Glu Cha Cys	0.3
17	Asp Glu Met Glu Nap Cys	0.8
18	AspS Val Val Abu Cys	4.6
19	Asp Glu Met Glu Glu Cys(Me)	16.7
20	Asp Glu Val Glu Cha Cys	0.33

21	Asp Glu Ile Glu Cha Cys	0.12
22	Asp Glu Tyr Glu Cha Cys	0.24
23	Asp Glu Phe Glu Cha Cys	0.42
24	Asp Glu Leu Glu Cha Cys	0.12
25	Asp Glu Cha Glu Cha Cys	0.14
26	Asp Glu Nle Glu Cha Cys	0.22
27	Asp Glu Tha Glu Cha Cys	0.87
28	Asp Glu FCI Glu Cha Cys	0.3
29	Asp Glu Phg Glu Cha Cys	0.12
30	Asp Glu Dif Glu Cha D-Cys	3.4
31	Glu Dif Glu Cha Cys	1.4
32	Dif Glu Cha Cys	30.0
33	Asp MGlu Leu Glu Cha Cys	1.0
34	Asp Glu Dif Glu Cha DHAla	7.1
35	Asp Glu Met Glu Glu Cpc	9.0
36	Glu Dif Ile Cha Cys	2.5
37	Dif Ile Cha Cys	100.0
38	Asp Glu Met Glu Glu CnAla	19.0
39	Asp Glu Leu Glu Cha Abu	1.6

40	Asp Glu Leu Glu Cha Val	4.0
41	Asp Glu Leu Glu Cha Nva	1.3
42	Asp-Asp-Leu-Glu-Cha-Cys	0.290
43	Asp-Fno-Leu-Glu-Cha-Cys	0.240
44	Asp-Tyr-Leu-Glu-Cha-Cys	0.135
45	Asp-(D)Phe-Leu-Glu-Cha-Cys	0.820
46	Asp-(D)Tyr-Leu-Glu-Cha-Cys	0.680
47	Asp-(D)Val-Leu-Glu-Cha-Cys	0.470
48	Asp-(D)Ile-Leu-Glu-Cha-Cys	0.330
49	Asp-(D)Dif-Leu-Glu-Cha-Cys	0,276
50	Asp-(D)Asp-Leu-Glu-Cha-Cys	0,122
51	Asp-Glu-Dap(N-b-Dns)-Glu-Cha-Cys	0,4

Particularly preferred examples of Pep, together with their IC_{50}s (in μM) are set out below in Table 3 are:

TABLE 3

Most preferred:

1	Asp Glu Dif Glu Cha Cys	0.05
2	Asp Glu Leu Val Cha Cys	0.08
3	Asp Glu Leu Ile Cha Cys	0.06
4	Asp Glu Dif Ile Cha Cys	0.06
5	Asp-Gla-Leu-Glu-Cha-Cys	0.055
6	Asp-(D)Glu-Leu-Glu-Cha-Cys	0.045
7	Asp-(D)Gla-Leu-Ile-Cha-Cys	0.0015
8	Glu-Leu-Glu-Cha-Cys	1.3
9	(D)Glu-Leu-Glu-Cha-Cys-(Pro-Cha-Asp-Leu)	0.080*
10	Succinyl Glu-Leu-Ile-Cha-Cys	
11	Succinyl (D)Glu-Leu-Glu-Cha-Cys-(Pro-Cha-Asp-Leu)	0.0040*
12	Asp-(D)Glu-Leu-Ile-Cha-Cys	
13	Asp-(D)Glu-Leu-Ile-Cha-Cys-(Pro-Cha-Asp-Leu)	<0.0002*

* Tested only as decapeptides

In these compounds:

Alg	=	allylglycine.
MGly	=	methylglycine.
MVal	=	methylvaline.
Abu	=	2-aminobutyric acid.
GluS	=	N-succinylglutamic acid.
AsGlu	=	Glutamic acid having N-terminal acylsulphonamide.
Cha	=	β-cyclohexylalanine.
Nap	=	naphthylalanine.
AspS	=	N-succinylaspartic acid.
Nle	=	norleucine.
Dif	=	3,3-diphenylalanine.
Tha	=	2-thienylalanine.
FCI	=	4-chlorophenylalanine.
Phg	=	phenylglycine.
CysMe	=	S-methylcysteine.

Cys(ACS)	=	Cysteine with C-terminal acylsulphonamide.
DHAla	=	dehydroalanine.
Cpc	=	1-amino-1-cyclopentane carboxylic acid.
CnAla	=	cyanoalanine.
MGlu	=	N-methylglutamic acid.
Fno	=	4-nitrophenylalanine.
Gla	=	γ-carboxyglutamic acid.
Dap	=	β-diaminopropionic acid.
Dns	=	dansyl(5-dimethylamino-1-naphthalene-sulfonyl).

Examples of compounds of the present invention may be effective as inhibitors of NS3 protease at micromolar or nanomolar levels. Preferably, the IC_{50}, as measured in the assay described below is less than 100nM, particularly preferably less than 20nM and, optimally, less than 5nM.

According to a second aspect, the present invention provides a compound, salt or derivative according to the first aspect, for use in any therapeutic method, preferably for use in inhibiting the HCV NS3 protease, and/or for use in treating or preventing hepatitis C or a related condition. By "related condition" is meant a condition which is or can be caused, directly or indirectly, by the hepatitis C virus, or with which the HCV is in any way associated.

According to a third aspect the present invention provides the use of a compound or derivative according to the first aspect in the manufacture of a medicament for the treatment or prevention of hepatitis C or a related condition.

A fourth aspect of the invention provides a pharmaceutical composition which includes one or more compounds or derivatives according to the first aspect.

The composition may also include pharmaceutically acceptable adjuvants such as carriers, buffers, stabilisers and other excipients. It may additionally include other therapeutically active agents, in particular those of use in treating or preventing hepatitis C or related conditions.

The pharmaceutical composition may be in any suitable form, depending on the intended method of administration. It may for example be in the form of a tablet, capsule or liquid for oral administration, or of a solution or suspension for administration parenterally.

According to a fifth aspect of the invention, there is provided a method of inhibiting HCV NS3 protease activity, and/or of treating or preventing hepatitis C or a related condition, the method involving administering to a human or animal (preferably mammalian) subject suffering from the condition a therapeutically or

prophylactically effective amount of a composition according to the fourth aspect of the invention, or of a compound or derivative according to the first aspect. "Effective amount" means an amount sufficient to cause a benefit to the subject or at least to cause a change in the subject's condition.

The dosage rate at which the compound, derivative or composition is administered will depend on the nature of the subject, the nature and severity of the condition, the administration method used, etc. Appropriate values can be selected by the trained medical practitioner. Preferred daily doses of the compounds are likely to be of the order of about 1 to 100 mg. The compound, derivative or composition may be administered alone or in combination with other treatments, either simultaneously or sequentially. It may be administered by any suitable route, including orally, intravenously, cutaneously, subcutaneously, etc. Intravenous administration is preferred. It may be administered directly to a suitable

site or in a manner in which it targets a particular site, such as a certain type of cell - suitable targeting methods are already known.

A sixth aspect of the invention provides a method of preparation of a pharmaceutical composition, involving admixing one or more compounds or derivatives according to the first aspect of the invention with one or more pharmaceutically acceptable adjuvants, and/or with one or more other therapeutically or prophylactically active agents.

According to a seventh aspect of the invention there is provided a method of producing the compounds of formula I. These compounds may be generated wholly or partly by chemical synthesis beginning from individual, preferably protected, amino acids or oligopeptides and using known peptide synthesis methods.

Modes for Carrying Out the Invention

Embodiments of the invention are exemplified below by way of illustration only.

EXAMPLES

(1) Synthesis

The synthesis of one of the compounds of the present invention is described below. Other compounds may be synthesized by an analogous method.

Synthesis of Ac-Asp-(D)Glu-Leu-Ile-Cha-Cys-Pro-Cha-Asp-Leu-Pro-Tyr-Lys(N^e-Ac)-NH_2

The synthesis was performed on solid phase by the continuous-flow Fmoc-polyamide method (Atherton, E. and Sheppard, R. C. (1989) Solid phase peptide synthesis. A practical approach, IRL Press, Oxford.). The resin used was Tentagel" derivatised with a modified Rink amide

linker p-[(R,S)-α-[1-(9H-Fluoren-9-yl)-methoxyformamido]-2,4-dimethoxybenzyl]-phenoxyacetic acid (Rink, H. (1987) *Tetrahedron Lett.* **28**, 3787-3789; Bernatowicz, M. S., Daniels, S. B. and Koster, H. (1989) *Tetrahedron Lett.* **30**, 4645-4667). All the coupling reactions were performed for 30 min with 5-fold excess of activated amino acid over the resin free amino groups, using Fmoc-amino acid/PyBOP/HOBt/DIEA (1:1:1:2) activation; double coupling was used for the cysteine residue. At the end of the assembly, the dry peptide-resin was treated with trifluoroacetic acid/water/triisopropylsilane (92.5:5:2.5) for 1.5h at room temperature; the resin was filtered out and the peptide precipitated with cold methyl t-Bu ether; the precipitate was redissolved in 50% water/acetonitrile containing 0.1%TFA and lyophillised.

Purification to >98% homogeneity was achieved through preparative HPLC on a Waters RCM (C-18) column (100 X 25 mm, 15mm) using as eluents (A) 0.1% trifluoroacetic acid in water and (B) 0.1%

trifluoroacetic acid in acetonitrile. The gradient used was 40%B isocratic for 5 min, then 40-60%B over 20 min, flow rate 30 ml/min; the fractions were analysed by HPLC (column: Beckman Ultrasphere, C-18, 25 X 4.6 mm, 5mm; gradient: 35-65%B in 20 min, same eluents as the preparative run, flow 1ml/min) and those containing the pure material were pooled and lyophilised (yield=50%). The Mass spectrum was acquired on a Perkin-Elmer API-100 spectrometer: MS= 1695.03 (calc.) 1694.6 (found).

(2) Inhibition Assay

The ability of the compounds to inhibit NS3 protease was evaluated using an NS3/4A complex comprising the NS3 protease domain and a modified form of the NS4A peptide, Pep 4AK [KKKGSVVIVGRIILSGR(NH_2)]. As substrate, a substrate peptide 4AB [DEMEECASHLPYK] based on the sequence of the NS4A/NS4B cleavage site of the HCV polyprotein, was used.

Cleavage assays were performed in 57µl 50 mM Hepes pH7.5, 1 % CHAPS, 15 % glycerol, 10 mM DTT (buffer A), to which 3µl substrate peptide were added. As protease co-factor a peptide spanning the central hydrophobic core (residues 21-34) of the NS4A protein, Pep4AK [KKKGSVVIVGRIILSGR(NH_2)] was used. Buffer solutions containing 80 µM Pep4AK were preincubated for 10 minutes with 10-200 nM protease and reactions were started by addition of substrate. Six duplicate data points at different substrate concentrations were used to calculate kinetic parameters. Incubation times were chosen in order to obtain <7% substrate conversion and reactions were stopped by addition of 40 µl 1 % TFA. Cleavage of peptide substrates was determined by HPLC using a Merck-Hitachi chromatograph equipped with an autosampler. 80 µl samples were injected on a Lichrospher C18 reversed phase cartridge column (4 x 74mm, 5µm, Merck) and fragments were separated using a 10-40 % acetonitrile gradient a 5%/min using a flow rate of 2.5ml/min. Peak detection was accomplished by monitoring both the

absorbance at 220nm and tyrosine fluorescence (λ_{ex} = 260 nm, λ_{em} = 305nm). Cleavage products were quantitated by integration of chromatograms with respect to appropriate standards. Kinetic parameters were calculated from nonlinear least-squares fit of initial rates as a function of substrate concentration with the help of a Kaleidagraph software, assuming Michaelis-Menten kinetics.

K_1 values of peptide inhibitors were calculated from substrate titration experiments performed in the presence of increasing amounts of inhibitor. Experimental data sets were simultaneously fitted to eq.1 using a multicurve fit macro with the help of a Sigmaplot software:

$$V = (V_{max}S)/(K_m(1+K_i/I)+S); \qquad \text{(eq.1)}$$

Alternatively, K_i values were derived from IC50

values, calculated using a four-parameter logistic function, according to eq.2:

$$IC50 - (1+S/K_m)K_i \qquad \text{(eq.2)}$$

The table below sets out the IC_{50} values for a variety of peptides tested in this assay and establishes that several optimised compounds of the present invention are active at nanomolar or subnanomolar levels. All the compounds tested - except for compound 26 which has a succinyl residue at the N-terminus- were tested as their N-acetyl derivatives.

Some of these compounds are the most potent in vitro inhibitors of HCV protease described to date. They are reversible, non covalent inhibitors which do not contain an electrophilic ("serine-trap") moiety in the molecule. They bind to both the S and S' region of the enzyme, and this makes them suitable for developing competition

binding assays, since they would be competitive with compounds binding to either the S or the S′ region of the enzyme.

Table 4

Ex. No	Sequence	IC_{50} (nM)
1	Glu Asp Val Val Abu Cys Pro Nle Ser Tyr	8500
2	Glu Asp Val Val Abu Cys (Me)Ala Nle Ser Tyr	3500
3	Asp Glu Dif Ile Cha Abu Ala Ser His Leu	29000
4	Asp Glu Dif Ile Cha Abu (Me)Ala Ser His Leu	29000
5	Asp Glu Dif Ile Cha (Me)Abu Ala Ser His Leu	8000
6	Asp Glu Dif Ile Cha (Me)Abu (Me)Ala Ser His Leu	3800
7	Asp (D)Glu Dif Ile Cha (Me)Abu (Me)Ala Ser His Leu	3100
8	Asp (D)Glu Leu Ile Cha Abu (Me)Ala Ser His Leu	5000
9	Asp Glu Dif Ile Cha Cys Pro Nle Ser Tyr	876
10	Glu Dif Ile Cha Cys Pro Nle Ser Leu	64
11	Asp Glu Dif Ile Cha Cys Pro Cha Ser Leu	23
12	Asp Glu Dif Ile Cha Cys Pro Cha Asp Leu	1.3
13	Asp Glu Dif Ile Cha Cys Pro Phg Asp Leu	7
14	Asp Glu Dif Ile Cha Cys Pro Nle Asp Leu	1.8
15	Asp Glu Dif Ile Cha Cys Pro Hof Asp Leu	1.8
16	Asp Glu Dif Ile Cha Cys Pro Cha Gln Leu	14
17	Asp Glu Dif Ile Cha Cys Pro Nle Gln Leu	32
18	Asp Glu Dif Ile Cha Cys Pro Hof Gln Leu	18

19	Asp Glu Dif Ile Cha Cys Pro Cha Hyp Leu	11
20	Asp Glu Dif Ile Cha Cys Pro Nle Hyp Leu	26
21	Asp Glu Dif Ile Cha Cys Pro Hof Hyp Leu	15
22	Asp (D)Glu Leu Ile Cha Cys Pro Nle Ser Leu	10
23	Asp Glu Dif Ile Cha Cys Pro Cha Asp Leu PYK(Ac)	0.85
24	Asp (D)Glu Leu Ile Cha Cys Pro Cha Asp Leu PYK(Ac)	< 0.2
25	Asp (D)Glu Leu Ile Cha Cys Pro Cha Asp Leu	< 0.2
26	Suc-(D)Glu Leu Ile Cha Cys Pro Cha Asp Leu	4
27	Asp (D)Glu Leu Glu Cha Cys Pro Cha Asp Leu	0.63
28	(D)Glu Leu Glu Cha Cys Pro Cha Asp Leu	80
29	Asp (D)Glu Leu Glu Cha Ala Pro Cha Asp Leu	17
30	Asp (D)Glu Leu Ile Cha Cys Pro Nle Ser Leu	10

Abbreviations used in Table I:

Abu = aminobutyric acid

Cha = β-cyclohexylalanine

Hof = homophenylalanine

Hyp = hydroxyproline

Lys(Ac) or K(Ac) = Nε-Acetyl-Lysine

Nle = norleucine

Phg = phenylglycine

Sta = statine [(3S,4S)-4-amino-3-hydroxy-6-methylheptanoic acid]

Dif = 3,3-diphenylalanine

Suc=succinyl

N-methylation is indicated as (Me) preceding the three-letter code of the amino acid

PYK = proline-tyrosine-lysine

(3) Substrate Assay

In order to determine whether or not an inhibitor molecule was a substrate for HCV NS3 protease a modified

version of the cleavage assay described above was employed using, as before, an NS3/4A complex comprising the NS3 protease domain and a modified form of the NS4A peptide, Pep4AK [KKKGSVVIVGRIILSGR(NH_2)].

1μM of the enzyme complex was incubated for 16hrs in the presence of 10μM inhibitor as a candidate substrate peptide. Assays were performed in 57μl 50 mM Hepes pH7.5, 1% CHAPS, 15% glycerol, 10 mM DTT.

After this time HPLC was used to separate any peptides resulting from cleavage and separated cleavage products detected.

Samples were analysed by HPLC on a Beckman 0.46 x 25 cm C18 reversed phase column equilibrated in 95% solvent A (0.1% TFA in H20) and 5% solvent B (0.1% TFA in acetonitrile) at a flow rate of 1 ml/min. Samples were eluted from this column with a linear gradient from 5% to

90% of B in 45 minutes. Peak detection was accomplished by monitoring absorbance at 220 nm.

CLAIMS:

1. A compound of formula (I), or a pharmaceutically acceptable salt or derivative thereof:

Pep-A'-B'-C'-D' (I)

wherein formula (I) is written from N-terminus on the left to C-terminus at the right and:

"Pep" is a peptide or peptide analogue capable of binding to HCV NS3 protease;

A' is a proline residue which is optionally substituted;

B' is an amino acid or amino acid analogue having a non-polar side chain;

C' is an amino acid or amino acid analogue having a polar side chain; and

D' is selected from leucine, other amino acids or amino acid analogues having a non-polar aliphatic side chain, and peptides of 2 to 6 amino acids having leucine or other amino acid or amino acid analogue with a non-polar aliphatic side chain as N-terminal residue;

and wherein the bond between Pep and A' is substantially uncleavable by HCV NS3 protease.

2. The compound of claim 1, a pharmaceutically acceptable salt or derivative thereof which is C-

terminally amidated.

3. The compound of claim 1 or claim 2, a pharmaceutically acceptable salt or derivative thereof which is N-terminally acylated.

4. A compound, salt or derivative according to any one of claims 1 to 3 wherein A'-B'-C'-D' is a tetrapeptide of formula:

Pro-B'-C'-Leu

wherein B' and C' are as defined in claim 1.

5. A compound, salt or derivative according to any one of the previous claims wherein B' is selected from: β-cyclohexylalanine, phenylglycine, homophenylalanine, norleucine, leucine, methionine, norvaline and β-cyclopropylalanine.

6. A compound, salt or derivative according to claim 5 wherein B' is selected from cyclohexylalanine and phenylglycine.

7. A compound, salt or derivative according to any one of the previous claims wherein C' is selected from: aspartic acid, glutamic acid, γ-carboxyglutamic acid,

glutamine, asparagine, hydroxyproline, N-β-Aloc-diaminobutyric acid, thiazolylalanine, methionine sulfoxide, pyridylalanine and serine.

8. A compound, salt or derivative according to claim 7 wherein C' is aspartic acid.

9. A compound, salt or derivative according to any one of the preceding claims, wherein the combination of amino acids B'C' is selected from:

Cha-Ser
Cha-Asp
Nle-Asp
Hof-Asp
Phg-Asp
Cha-Gln
Nle-Gln
Hof-Gln
Cha-Hyp
Nle-Hyp
Hof-Hyp
Nle-Ser

10. A compound, salt, or derivative according to any one of the preceding claims wherein Pep-OH is capable of

binding HCV NS3 protease, in the absence of the C-terminal residues A'-B'-C'-D', and has an IC_{50} below 100μM in an inhibition assay.

11. A compound, salt, or derivative according to any one of the preceding claims wherein Pep is a hexa-, penta- or tetra-peptide having formula (II) below:

F-E-D-C-B-A

wherein: A is an amino acid or amino acid analogue having an aliphatic side chain of form 1 to 6 carbon atoms;

B is an amino acid or analogue having a non-polar, acidic, or polar but uncharged side group;

C is an amino acid or amino acid analogue having a non-polar or acidic side chain;

D is an amino acid or amino acid analogue having a hydrophobic side group;

E together with F may be absent, but if present is an amino acid or amino acid analogue having an acidic side chain, non-polar side chain or polar, but uncharged side chain, or is a dicarboxylic acid containing up to 6 carbon atoms and lacking the amino group of acidic amino acids;

and F may be absent (either by itself, or together with E) but when present is an amino acid or analogue

having an acidic side chain or is a dicarboxylic acid containing up to 6 carbon atoms.

12. A compound according to claim 11 wherein:

A is selected from: cysteine, aminobutyric acid, di- and tri-fluoro aminobutyric acid, norvaline, allylglycine and alanine;

B is selected from: glutamic acid, aspartic acid, glycine, methyl glycine, 2-amino butyric acid, alanine, isoleucine, valine, leucine, cysteine, naphthylalanine and β-cyclohexylalanine;

C is selected from: glutamic acid, aspartic acid, glycine, methyl glycine, 2-amino butyric acid, alanine, isoleucine, valine, leucine, cysteine, naphthylalanine and β-cyclohexylalanine;

D is selected from: methionine, isoleucine, leucine, norleucine, valine, methylvaline, phenylglycine, diphenylalanine, tyrosine, thienylalanine, and chlorophenylalanine;

E is selected from: glutamic acid, aspartic acid, phenylalanine, diphenylalanine, isoleucine, valine, tyrosine, 4-nitrophenylalanine, glutaric acid and succinic acid;

and F is selected from: aspartic acid, glutamic acid, glutaric acid and succinic acid.

13. A compound, salt, or derivative, according to any one of the preceding claims for use in therapy.

14. A pharmaceutical composition comprising a compound, salt or derivative according to any one of the preceding claims and a pharmaceutically acceptable excipient, diluent or carrier.

15. Use of a compound, salt or derivative according to any one of the preceding claims in the manufacture of a medicament for the treatment or prevention of hepatitis C or a related condition.

16. A method of inhibiting HCV NS3 protease activity, and/or of treating or preventing hepatitis C or a related condition, comprising administering to a human or mammalian subject suffering from the condition a therapeutically or prophylactically effective amount of a composition according to claim 14, or of a compound of any one of claims 1 to 12.

09856886 .072604

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 :
C07K 14/18, A61K 39/29, A61P 31/20

A1

(11) International Publication Number: WO 00/31129

(43) International Publication Date: 2 June 2000 (02.06.00)

(21) International Application Number: PCT/EP99/09207

(22) International Filing Date: 24 November 1999 (24.11.99)

(30) Priority Data:
9825946.8 26 November 1998 (26.11.98) GB

(71) Applicant *(for all designated States except US)*: ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P ANGELETTI SPA [IT/IT]; Via Pontina KM 30.600, I–00040 Pomezia (IT).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: PESSI, Antonello [IT/IT]; Via Monte Bianco, 63, I–00141 Rome (IT). INGALLINELLA, Paola [IT/IT]; Via Zara, 15, I–00040 Pomezia (IT). BIANCHI, Elisabetta [IT/IT]; Via Sabotino, 31, I–00195 Rome (IT).

(74) Agents: NICHOLLS, Kathryn, M. et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: PHARMACEUTICAL COMPOUNDS FOR THE INHIBITION OF HEPATITIS C VIRUS NS3 PROTEASE

(57) Abstract

Peptidic inhibitors of hepatitis C virus NS3 protease are disclosed which are based on the P and P' regions of the natural substrate. The P' part of the inhibitor is optimised to achieve maximum binding energy through interaction with the S' region of the enzyme. By selecting amino acids such that the inhibitor is substantially not cleavable by the NS3 protease inhibitors having potency in the low nanomolar to sub–nanomolar range can be achieved.

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Includes Reference to PCT International Applications)	ATTORNEY'S DOCKET NUMBER IT0013

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Pharmaceutical compounds for the inhibition of hepatitis C virus NS3 protease

the specification of which (check only one item below):

☐ is attached hereto.

☐ was filed as United States application

Serial No.

on

and was amended

on (if applicable).

☒ was filed as PCT international application

Number PCT/EP99/09207

on 24 November 1999

and was amended under PCT Article 19

on (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:

COUNTRY (if PCT indicate PCT)	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
Great Britain	9825946.8	26 November 1998	☒ YES ☐ NO
			☐ YES ☐ NO
			☐ YES ☐ NO
			☐ YES ☐ NO
			☐ YES ☐ NO

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Continued) (includes Reference to PCT International Applications)	ATTORNEY'S DOCKET NUMBER IT0013

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:

U.S. APPLICATIONS		STATUS *(Check one)*		
U S APPLICATION NUMBER	U S FILING DATE	PATENTED	PENDING	ABANDONED
PCT APPLICATIONS DESIGNATING THE U.S.				
PCT APPLICATION NO	PCT FILING DATE / US SERIAL NUMBERS ASSIGNED *(if any)*			
		☐	☐	☐

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(List name and registration number)* *S. O. Heber - 38,179 J. L. Tribble - 32,633*

Send Correspondence to Merck & Co., Inc., 126 E. Lincoln Avenue, Rahway, New Jersey 07065, U.S.A.				Direct Telephone Calls to *(name and telephone number)* *S. O. Heber (732)594-1958*
2	FULL NAME OF INVENTOR	FAMILY NAME Bianchi	FIRST GIVEN NAME Elisabetta	SECOND GIVEN NAME
0	RESIDENCE & CITIZENSHIP	CITY Rome	STATE OR FOREIGN COUNTRY Italy	COUNTRY OF CITIZENSHIP Italy
1	POST OFFICE ADDRESS	POST OFFICE ADDRESS IRBM, Via Pontina KM 30.600	CITY Pomezia (Rome)	STATE & ZIP CODE/COUNTRY I-00040, Italy
2	FULL NAME OF INVENTOR	FAMILY NAME Ingallinella	FIRST GIVEN NAME Paolo	SECOND GIVEN NAME
0	RESIDENCE & CITIZENSHIP	CITY Naples	STATE OR FOREIGN COUNTRY Italy	COUNTRY OF CITIZENSHIP Italy
2	POST OFFICE ADDRESS	POST OFFICE ADDRESS IRBM, Via Pontina KM 30.600	CITY Pomezia (Rome)	STATE & ZIP CODE/COUNTRY I-00040, Italy
2	FULL NAME OF INVENTOR	FAMILY NAME Pessi	FIRST GIVEN NAME Antonello	SECOND GIVEN NAME
0	RESIDENCE & CITIZENSHIP	CITY Rome	STATE OR FOREIGN COUNTRY Italy	COUNTRY OF CITIZENSHIP Italy
3	POST OFFICE ADDRESS	POST OFFICE ADDRESS IRBM, Via Pontina KM 30.600	CITY Pomezia (Rome)	STATE & ZIP CODE/COUNTRY I-00040, Italy

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201	SIGNATURE OF INVENTOR 202	SIGNATURE OF INVENTOR 203
DATE	DATE May 31st 2001	DATE

09856886.072601

Rec'd PCT/PTO 17 OCT 2002

IT0013P



SEQUENCE LISTING

<110> Bianchi, Elisabetta
Ingallinella, Paolo
Pessi, Antonello

<120> PHARMACEUTICAL COMPOUNDS FOR THE INHIBITION OF HEPATITIS C VIRUS NS3 PROTEASE

<130> IT0013P

<140> 09/856,886
<141> 2001-07-26

<150> PCT/EP99/09207
<151> 1999-11-24

<150> GB/9825946.8
<151> 1998-11-26

<160> 109

<170> FastSEQ for Windows Version 4.0

<210> 1
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 2
<223> Xaa=beta-cyclohexylalanine

<400> 1
Pro Xaa Ser Leu
1

<210> 2
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 2
<223> Xaa=beta-cyclohexylalanine

<400> 2
Pro Xaa Asp Leu
1

<210> 3
<211> 4
<212> PRT

<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 2
<223> Nle

<400> 3
Pro Leu Asp Leu
 1

<210> 4
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 2
<223> Xaa=homophenylalanine

<400> 4
Pro Xaa Asp Leu
 1

<210> 5
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 2
<223> Xaa=phenylglycine

<400> 5
Pro Xaa Asp Leu
 1

<210> 6
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 2
<223> Xaa=beta-cyclohexylalanine

<400> 6

Pro Xaa Gln Leu
1

<210> 7
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 2
<223> Nle

<400> 7
Pro Leu Gln Leu
1

<210> 8
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 2
<223> Xaa=homophenylalanine

<400> 8
Pro Xaa Gln Leu
1

<210> 9
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 2
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 3
<223> Xaa=hydroxyproline

<400> 9
Pro Xaa Xaa Leu
1

<210> 10
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 2
<223> Nle

<221> SITE
<222> 3
<223> Xaa=hydroxyproline

<400> 10
Pro Leu Xaa Leu
 1

<210> 11
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 2
<223> Xaa=homophenylalanine

<221> SITE
<222> 3
<223> Xaa=hydroxyproline

<400> 11
Pro Xaa Xaa Leu
 1

<210> 12
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 2
<223> Nle

<400> 12
Pro Leu Ser Leu
 1

<210> 13
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<400> 13
Asp Glu Met Glu Glu Cys
 1 5

<210> 14
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 6
<223> Xaa=D-Cys

<400> 14
Asp Glu Met Glu Glu Xaa
 1 5

<210> 15
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 6
<223> Xaa=Abu

<400> 15
Asp Glu Met Glu Glu Xaa
 1 5

<210> 16
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1

<223> Acetylation

<400> 16
Met Glu Glu Cys
1

<210> 17
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<400> 17
Glu Met Glu Glu Cys
1 5

<210> 18
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<400> 18
Glu Asp Val Val Cys Cys
1 5

<210> 19
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=Abu

<400> 19
Glu Asp Val Val Xaa Cys
1 5

<210> 20

0956396.072601

<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<400> 20
Asp Glu Val Val Cys Cys
 1 5

<210> 21
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<400> 21
Glu Asp Val Val Gly Cys
 1 5

<210> 22
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 6
<223> Xaa=allylglycine

<400> 22
Asp Glu Met Glu Glu Xaa
 1 5

<210> 23
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> MOD_RES
<222> 5
<223> MeGly

<400> 23
Glu Asp Val Val Gly Cys
 1 5

<210> 24
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> MOD_RES
<222> 3
<223> MeVal

<221> SITE
<222> 5
<223> Xaa=Abu

<400> 24
Glu Asp Val Val Xaa Cys
 1 5

<210> 25
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 1
<223> Xaa=N-succinylglutamic acid

<400> 25
Xaa Met Glu Glu Cys
 1 5

<210> 26
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 1
<223> Xaa=glutamic acid having N-terminal
acylsulphonamide

<400> 26
Xaa Met Glu Glu Cys
1 5

<210> 27
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<400> 27
Asp Glu Met Glu Leu Cys
1 5

<210> 28
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 28
Asp Glu Met Glu Xaa Cys
1 5

<210> 29
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE

<222> 5
<223> Xaa=napthylalanine

<400> 29
Asp Glu Met Glu Xaa Cys
1 5

<210> 30
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 1
<223> Xaa=N-succinylaspartic acid

<221> SITE
<222> 4
<223> Xaa=Abu

<400> 30
Xaa Val Val Xaa Cys
1 5

<210> 31
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 6
<223> Xaa=S-methylcysteine

<400> 31
Asp Glu Met Glu Glu Xaa
1 5

<210> 32
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 32
Asp Glu Val Glu Xaa Cys
 1 5

<210> 33
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 33
Asp Glu Ile Glu Xaa Cys
 1 5

<210> 34
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 34
Asp Glu Tyr Glu Xaa Cys
 1 5

<210> 35
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 35
Asp Glu Phe Glu Xaa Cys
 1 5

<210> 36
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 36
Asp Glu Leu Glu Xaa Cys
 1 5

<210> 37
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 37
Asp Glu Xaa Glu Xaa Cys
 1 5

<210> 38
<211> 6
<212> PRT
<213> Artificial Sequence

<220>

<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> MOD_RES
<222> 3
<223> Nle

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 38
Asp Glu Leu Glu Xaa Cys
 1 5

<210> 39
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=2-thienylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 39
Asp Glu Xaa Glu Xaa Cys
 1 5

<210> 40
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=4-chlorophenylalanine

<221> SITE

<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 40
Asp Glu Xaa Glu Xaa Cys
 1 5

<210> 41
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=phenylglycine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 41
Asp Glu Xaa Glu Xaa Cys
 1 5

<210> 42
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 6
<223> Xaa=D-Cys

<400> 42
Asp Glu Xaa Glu Xaa Xaa
 1 5

<210> 43
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 4
<223> Xaa=beta-cyclohexylalanine

<400> 43
Glu Xaa Glu Xaa Cys
 1 5

<210> 44
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 1
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 3
<223> Xaa=beta-cyclohexylalanine

<400> 44
Xaa Glu Xaa Cys
 1

<210> 45
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=N-methylglutamic acid

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 45
Asp Xaa Leu Glu Xaa Cys
 1 5

<210> 46
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 6
<223> Xaa=dehydroalanine

<400> 46
Asp Glu Xaa Glu Xaa Xaa
 1 5

<210> 47
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 6
<223> Xaa=1-amino-1-cyclopentane carboxylic acid

<400> 47

0985686.072601

Asp Glu Met Glu Glu Xaa
1 5

<210> 48
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 4
<223> Xaa=beta-cyclohexylalanine

<400> 48
Glu Xaa Ile Xaa Cys
1 5

<210> 49
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 1
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 3
<223> Xaa=beta-cyclohexylalanine

<400> 49
Xaa Ile Xaa Cys
1

<210> 50
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 6
<223> Xaa=cyanoalanine

<400> 50
Asp Glu Met Glu Glu Xaa
 1 5

<210> 51
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 6
<223> Xaa=Abu

<400> 51
Asp Glu Leu Glu Xaa Xaa
 1 5

<210> 52
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 52
Asp Glu Leu Glu Xaa Val
 1 5

<210> 53
<211> 6
<212> PRT

<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 6
<223> Nva

<400> 53
Asp Glu Leu Glu Xaa Val
 1 5

<210> 54
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 54
Asp Asp Leu Glu Xaa Cys
 1 5

<210> 55
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=4-nitrophenylalanine

<221> SITE
<222> 5

0985886.072601

<223> Xaa=beta-cyclohexylalanine

<400> 55
Asp Xaa Leu Glu Xaa Cys
 1 5

<210> 56
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 56
Asp Tyr Leu Glu Xaa Cys
 1 5

<210> 57
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Phe

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 57
Asp Xaa Leu Glu Xaa Cys
 1 5

<210> 58
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES

<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Tyr

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 58
Asp Xaa Leu Glu Xaa Cys
 1 5

<210> 59
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Val

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 59
Asp Xaa Leu Glu Xaa Cys
 1 5

<210> 60
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Ile

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 60
Asp Xaa Leu Glu Xaa Cys
1 5

<210> 61
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 61
Asp Xaa Leu Glu Xaa Cys
1 5

<210> 62
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Asp

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 62
Asp Xaa Leu Glu Xaa Cys
1 5

<210> 63
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=beta-diaminopropionic acid

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 3
<223> N-b- dansyl
(5-dimethylamino-1-naphthalene-sulfonyl)

<400> 63
Asp Glu Xaa Glu Xaa Cys
1 5

<210> 64
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 64
Asp Glu Xaa Glu Xaa Cys
1 5

<210> 65
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE

<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 65
Asp Glu Leu Val Xaa Cys
 1 5

<210> 66
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 66
Asp Glu Leu Ile Xaa Cys
 1 5

<210> 67
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 67
Asp Glu Xaa Ile Xaa Cys
 1 5

<210> 68
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=gamma-carboxyglutamic acid

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 68
Asp Xaa Leu Glu Xaa Cys
 1 5

<210> 69
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 69
Asp Xaa Leu Glu Xaa Cys
 1 5

<210> 70
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)gamma-carboxyglutamic acid

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 70
Asp Xaa Leu Ile Xaa Cys
 1 5

<210> 71
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 4
<223> Xaa=beta-cyclohexylalanine

<400> 71
Glu Leu Glu Xaa Cys
 1 5

<210> 72
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 1
<223> Xaa=(D)Glu

<221> SITE
<222> 4
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 7
<223> Xaa=beta-cyclohexylalanine

<400> 72
Xaa Leu Glu Xaa Cys Pro Xaa Asp Leu
 1 5

<210> 73
<211> 5
<212> PRT
<213> Artificial Sequence

<220>

<223> Synthetic sequence

<221> SITE
<222> 4
<223> Xaa=beta-cyclohexylalanine

<400> 73
Glu Leu Ile Xaa Cys
 1 5

<210> 74
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 1
<223> Xaa=(D)Glu

<221> SITE
<222> 4, 7
<223> Xaa=beta-cyclohexylalanine

<400> 74
Xaa Leu Glu Xaa Cys Pro Xaa Asp Leu
 1 5

<210> 75
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<400> 75
Asp Xaa Leu Ile Xaa Cys
 1 5

<210> 76
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<400> 76
Asp Xaa Leu Ile Xaa Cys Pro Xaa Asp Leu
 1 5 10

<210> 77
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 13
<223> Ne-Acetyl-Lysine

<221> MOD_RES
<222> 13
<223> Amidation

<400> 77
Asp Xaa Leu Ile Xaa Cys Pro Xaa Asp Leu Pro Tyr Lys
 1 5 10

<210> 78
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<400> 78
Lys Lys Lys Gly Ser Val Val Ile Val Gly Arg Ile Ile Leu Ser Gly
 1 5 10 15
Arg

<210> 79
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<400> 79
Asp Glu Met Glu Glu Cys Ala Ser His Leu Pro Tyr Lys
 1 5 10

<210> 80
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=Abu

<221> MOD_RES
<222> 8
<223> Nle

<400> 80
Glu Asp Val Val Xaa Cys Pro Leu Ser Tyr
 1 5 10

<210> 81
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 5
<223> Xaa=Abu

<221> SITE
<222> 7

<223> N-methylation

<221> MOD_RES
<222> 8
<223> Nle

<400> 81
Glu Asp Val Val Xaa Cys Ala Leu Ser Tyr
 1 5 10

<210> 82
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 6
<223> Xaa=Abu

<400> 82
Asp Glu Xaa Ile Xaa Xaa Ala Ser His Leu
 1 5 10

<210> 83
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE

<222> 6
<223> Xaa=Abu

<221> SITE
<222> 7
<223> N-methylation

<400> 83
Asp Glu Xaa Ile Xaa Xaa Ala Ser His Leu
 1 5 10

<210> 84
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 6
<223> Xaa=Abu

<221> SITE
<222> 6
<223> Xaa=N-methylation

<400> 84
Asp Glu Xaa Ile Xaa Xaa Ala Ser His Leu
 1 5 10

<210> 85
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 6
<223> Xaa=Abu

<221> SITE
<222> 6
<223> Xaa=N-methylation

<400> 85
Asp Glu Xaa Ile Xaa Xaa Ala Ser His Leu
 1 5 10

<210> 86
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 6
<223> Xaa=Abu

<221> SITE
<222> (6)...(7)
<223> N-methylation

<400> 86
Asp Xaa Xaa Ile Xaa Xaa Ala Ser His Leu
 1 5 10

<210> 87
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 6
<223> Xaa=Abu

<221> SITE
<222> 7
<223> N-methylation

<400> 87
Asp Xaa Leu Ile Xaa Xaa Ala Ser His Leu
 1 5 10

<210> 88
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 8
<223> Nle

<400> 88
Asp Glu Xaa Ile Xaa Cys Pro Leu Ser Tyr
 1 5 10

<210> 89
<211> 9
<212> PRT
<213> Artificial Sequence

<220>

<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 4
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 7
<223> Nle

<400> 89
Glu Xaa Ile Xaa Cys Pro Leu Ser Leu
 1 5

<210> 90
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<400> 90
Asp Glu Xaa Ile Xaa Cys Pro Xaa Ser Leu
 1 5 10

<210> 91
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE

<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<400> 91
Asp Glu Xaa Ile Xaa Cys Pro Xaa Asp Leu
 1 5 10

<210> 92
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 8
<223> Xaa=phenylglycine

<400> 92
Asp Glu Xaa Ile Xaa Cys Pro Xaa Asp Leu
 1 5 10

<210> 93
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 8
<223> Nle

<400> 93
Asp Glu Xaa Ile Xaa Cys Pro Leu Asp Leu
 1 5 10

<210> 94
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 8
<223> Xaa=homophenylalanine

<400> 94
Asp Glu Xaa Ile Xaa Cys Pro Xaa Asp Leu
 1 5 10

<210> 95
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<400> 95
Asp Glu Xaa Ile Xaa Cys Pro Xaa Gln Leu
 1 5 10

<210> 96
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 8
<223> Nle

<400> 96
Asp Glu Xaa Ile Xaa Cys Pro Leu Gln Leu
 1 5 10

<210> 97
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 8
<223> Xaa=homophenylalanine

<400> 97
Asp Glu Xaa Ile Xaa Cys Pro Xaa Gln Leu
 1 5 10

<210> 98
<211> 10

<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> SITE
<222> 8
<223> Xaa=homophenylalanine

<221> SITE
<222> 9
<223> Xaa=hydroxyproline

<400> 100
Asp Glu Xaa Ile Xaa Cys Pro Xaa Xaa Leu
 1 5 10

<210> 101
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 8
<223> Nle

<400> 101
Asp Xaa Leu Ile Xaa Cys Pro Leu Ser Leu
 1 5 10

<210> 102
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 3
<223> Xaa=3,3-diphenylalanine

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 13
<223> Ne-Acetyl-Lysine

<400> 102
Asp Glu Xaa Ile Xaa Cys Pro Xaa Asp Leu Pro Tyr Lys
 1 5 10

<210> 103
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 13
<223> Ne-Acetyl-Lysine

<400> 103
Asp Xaa Leu Ile Xaa Cys Pro Xaa Asp Leu Pro Tyr Lys
 1 5 10

<210> 104
<211> 10

<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)GLu

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<400> 104
Asp Xaa Leu Ile Xaa Cys Pro Xaa Asp Leu
 1 5 10

<210> 105
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> SITE
<222> 1
<223> Xaa=(D)Glu

<221> SITE
<222> 4, 7
<223> Xaa=beta-cyclohexylalanine

<400> 105
Xaa Leu Ile Xaa Cys Pro Xaa Asp Leu
 1 5

<210> 106
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE

<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<400> 106
Asp Xaa Leu Glu Xaa Cys Pro Xaa Asp Leu
 1 5 10

<210> 107
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 1
<223> Xaa=(D)Glu

<221> SITE
<222> 4, 7
<223> Xaa=beta-cyclohexylalanine

<400> 107
Xaa Leu Glu Xaa Cys Pro Xaa Asp Leu
 1 5

<210> 108
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5, 8
<223> Xaa=beta-cyclohexylalanine

<400> 108
Asp Xaa Leu Glu Xaa Ala Pro Xaa Asp Leu
 1 5 10

<210> 109
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic sequence

<221> MOD_RES
<222> 1
<223> Acetylation

<221> SITE
<222> 2
<223> Xaa=(D)Glu

<221> SITE
<222> 5
<223> Xaa=beta-cyclohexylalanine

<221> MOD_RES
<222> 8
<223> Nle

<400> 109
Asp Xaa Leu Ile Xaa Cys Pro Leu Ser Leu
 1 5 10

binding HCV NS3 protease, in the absence of the C-terminal residues A'-B'-C'-D', and has an IC_{50} below 100μM in an inhibition assay.

11. A compound, salt, or derivative according to any one of the preceding claims wherein Pep is a hexa-, penta- or tetra-peptide having formula (II) below:

F-E-D-C-B-A

wherein: A is an amino acid or amino acid analogue having an aliphatic side chain of form 1 to 6 carbon atoms;

B is an amino acid or analogue having a non-polar, acidic, or polar but uncharged side group;

C is an amino acid or amino acid analogue having a non-polar or acidic side chain;

D is an amino acid or amino acid analogue having a hydrophobic side group;

E together with F may be absent, but if present is an amino acid or amino acid analogue having an acidic side chain, non-polar side chain or polar, but uncharged side chain, or is a dicarboxylic acid containing up to 6 carbon atoms and lacking the amino group of acidic amino acids;

and F may be absent (either by itself, or together with E) but when present is an amino acid or analogue